

Understanding Breast Cancer

A guide for people with cancer, their families and friends



For information & support
call **Cancer Connect 13 11 20**

Understanding Breast Cancer

A guide for people with cancer, their families and friends

First published May 2006. This edition September 2026.

© Cancer Council Australia 2026. ISBN 978 1 923073 56 2

Understanding Breast Cancer is reviewed approximately every 3 years.

Check the publication date above to ensure this copy is up to date.

Editor: Gina Flaxman. Designer: Caen Clasby. Printer: IVE Group.

Acknowledgements

This edition has been developed by Cancer Council NSW on behalf of all other state and territory Cancer Councils as part of a National Cancer Information Subcommittee initiative. We thank the reviewers of this booklet: A/Prof Cindy Mak, Breast Surgeon and Director of Breast Services, Chris O'Brien Lifehouse, Royal Prince Alfred Hospital, NSW; A/Prof Nick Murray, Consultant Medical Oncologist, Royal Adelaide Hospital and GenesisCare Adelaide, SA; Caroline Bennett, Consumer; Tamara Casey, Clinical Nurse Consultant, Breast Assessment Unit, Fiona Stanley Hospital, WA; A/Prof Joe Dusseldorp, Reconstructive Plastic Surgeon, Chris O'Brien Lifehouse, NSW; Lauren Lores, Consumer; Dr Linda Madigan, Breast Physician, Royal Adelaide Hospital, SA; Catherine Meredith, Consumer; Ash Mondolo, Breast Cancer Clinical Nurse Consultant, Mater Private Hospital, QLD; Chris Rivett, 13 11 20 Consultant, Cancer Council SA; Catrina Ross, McGrath Breast Care Nurse, Northern Sydney Local Health District, NSW; Prof Christobel Saunders AO, James Stewart Chair of Surgery and Director of Research, Melbourne Medical School, The University of Melbourne, VIC; Dr Janice Yeh, Radiation Oncologist, Peter MacCallum Cancer Centre, VIC; A/Prof Yvonne Zissiadis, Radiation Oncologist, Medical Director, GenesisCare WA and Adjunct A/Prof, Curtin University, WA. We also thank the health professionals, consumers and editorial teams who have worked on previous editions of this title.

The photographs on page 37 have been reproduced with permission from (from top): A/Prof James French, Westmead Breast Cancer Institute, NSW; Prof Elisabeth Elder, Westmead Breast Cancer Institute, The University of Sydney, NSW; A/Prof Joe Dusseldorp, Reconstructive Plastic Surgeon, Chris O'Brien Lifehouse, NSW.

Note to reader

Always consult your doctor about matters that affect your health. This booklet is intended as a general introduction to the topic and should not be seen as a substitute for medical, legal or financial advice. You should obtain independent advice relevant to your specific situation from appropriate professionals, and you may wish to discuss issues raised in this booklet with them. All care is taken to ensure that the information in this booklet is accurate at the time of publication. Please note that information on cancer, including the diagnosis, treatment and prevention of cancer, is constantly being updated and revised by medical professionals and the research community. Cancer Council Australia and its members exclude all liability for any injury, loss or damage incurred by use of or reliance on the information provided in this booklet.

Cancer Council

Cancer Council is Australia's peak non-government cancer control organisation. Through the 8 state and territory Cancer Councils, we provide a broad range of programs and services to help improve the quality of life of people living with cancer, their families and friends. Cancer Councils also invest heavily in research and prevention. To make a donation and help us beat cancer, visit cancer.org.au or call your local Cancer Council.



Cancer Council acknowledges Traditional Custodians of Country throughout Australia and recognises the continuing connection to lands, waters and communities. We pay our respects to Aboriginal and Torres Strait Islander cultures and to Elders past and present.



Cancer Council Australia Suite 1202, Level 12, Centennial Place,
300 Elizabeth Street, Sydney NSW 2010 **Telephone** 02 8256 4100
Email info@cancer.org.au **Website** cancer.org.au **ABN** 91 130 793 725

About this booklet

This booklet has been prepared to help you understand more about early and locally advanced breast cancer. It also includes a short section on advanced (metastatic) breast cancer.

Many people feel shocked and upset when told they have breast cancer. We hope this booklet will help you, your family and friends understand how breast cancer is diagnosed and treated. We also include information about support services.

We cannot give advice about the best treatment for you. You need to discuss this with your doctors. However, this information may answer some of your questions and help you think about what to ask your treatment team (see page 67 for a question checklist).

This booklet does not need to be read from cover to cover – just read the parts that are useful to you. Some medical terms that may be unfamiliar are explained in the glossary (see page 68). You may also like to pass this booklet to family and friends for their information.

How this booklet was developed – This information was developed with help from a range of health professionals and people affected by breast cancer. It is based on Australian clinical practice guidelines for early breast cancer and international guidelines for advanced breast cancer.¹⁻²



If you or your family have any questions or concerns, call **Cancer Connect on 13 11 20**. We can send you more information and connect you with support services in your area. You can also visit the Cancer Connect website at cancerconnect.org.au.



Contents

What is cancer?	4
The breasts	6
Key questions	8
What is breast cancer?	8
How common is breast cancer?	8
Does breast cancer run in families?	9
What are the risk factors for breast cancer?	10
What are the different types of breast conditions and breast cancers?	12
What are the symptoms?	13
Which health professionals will I see?	13
Diagnosis	16
Mammogram	16
Ultrasound	17
Breast MRI scan	18
Biopsy	18
Further tests	19
Staging breast cancer	20
Grading breast cancer	21
Prognosis	21
Making treatment decisions	25
Treatment	27
Surgery	27
Radiation therapy	40
Chemotherapy	43
Hormone therapy	45

Targeted therapy	47
------------------	----

Immunotherapy	49
---------------	----

Managing side effects	52
------------------------------	-----------

Lymphoedema	52
-------------	----

Cording	53
---------	----

Nerve pain	53
------------	----

Hair loss	54
-----------	----

Thinking and memory changes	54
-----------------------------	----

Breast prostheses	55
-------------------	----

Changes to body image and sex life	56
------------------------------------	----

Menopause and infertility	57
---------------------------	----

Looking after yourself	58
-------------------------------	-----------

Life after treatment	60
-----------------------------	-----------

Follow-up appointments	61
------------------------	----

What if the cancer returns?	62
-----------------------------	----

Caring for someone with cancer	63
---------------------------------------	-----------

Seeking support	64
------------------------	-----------

Support from Cancer Council	65
-----------------------------	----

Useful websites	66
-----------------	----

Question checklist	67
---------------------------	-----------

Glossary	68
-----------------	-----------

How you can help	72
-------------------------	-----------

Key to icons

Icons are used throughout this booklet to indicate:



More information



Alert



Tips

Is this Cancer Council booklet helpful?

Please follow this QR code for a quick 3-minute survey, or call 13 11 20 to provide your feedback.



SCAN ME


What is cancer?

Cancer is a disease of the cells. Cells are the body's basic building blocks – they make up tissues and organs. The body constantly makes new cells to help us grow, replace worn-out tissue and heal injuries.

Normally, cells multiply and die in an orderly way, so that each new cell replaces one lost. Sometimes, however, cells become abnormal and keep growing. These abnormal cells may turn into cancer.

In solid cancers, such as breast cancer, the abnormal cells form a mass or lump called a tumour. In some cancers, such as leukaemia, the abnormal cells build up in the blood.

How cancer starts





Not all tumours are cancer. Benign tumours tend to grow slowly and usually don't move into other parts of the body or turn into cancer. Cancerous tumours, also known as malignant tumours, have the potential to spread. They may invade nearby tissue, destroying normal cells. The cancer cells can break away and travel through the bloodstream or lymph vessels to other parts of the body.

The cancer that first develops in a tissue or organ is called the primary cancer. It is considered localised cancer if it has not spread to other parts of the body. If the primary cancer cells grow and form another tumour at a new site, it is called a secondary cancer or metastasis. A metastasis keeps the name of the original cancer. For example, breast cancer that has spread to the liver is called metastatic breast cancer, even though the main symptoms may be coming from the liver.

How cancer spreads

.....► Malignant cancer



The breasts

The breasts sit on top of the upper ribs and a large chest muscle. They cover the area from the collarbone (clavicle) to the armpit (axilla) and across to the breastbone (sternum). Some breast tissue extends into the armpit and is called the axillary tail. Female breasts are mostly made up of:


- **lobes** – each breast has 12-20 sections called lobes
- **lobules** – each lobe contains glands that can produce milk; these milk glands are called lobules or glandular tissue
- **ducts** – the lobes and lobules are connected by fine tubes called ducts; the ducts carry milk to the nipples when breastfeeding
- **fatty/fibrous tissue** – all breasts contain some fatty and fibrous tissue (including connective tissue called stroma), no matter their size.

Most younger women have dense or thicker breasts because they contain more lobules than fat. Male breasts have ducts and fatty/fibrous tissue. They usually contain no, or only a few, lobes and lobules.

The lymphatic system – The lymphatic system is an important part of the immune system, which protects against disease and infection. The lymphatic system drains excess fluid from body tissues into the blood. It is made up of a network of thin tubes called lymph vessels. These vessels connect to groups of small, bean-shaped lymph nodes (or glands).

There are lymph nodes throughout the body, including in the armpits, neck, abdomen, groin and chest (near the breastbone). The first place breast cancer cells usually spread to is the lymph nodes in the armpits (axillary lymph nodes) or to the lymph nodes near the breastbone (internal mammary lymph nodes).

The breasts



Everyone has breast tissue, regardless of their gender. We understand that not everyone identifies with having “breasts” and that some people find the word “breast” a gendered word and prefer the word “chest”. We use the term “breast” to refer to the tissue affected by breast cancer (breast tissue). Chest cancers are thoracic cancers like lung and tracheal (windpipe) cancer.

Key questions

Q: What is breast cancer?

A: Breast cancer usually begins when cells in the breast become abnormal. The cells grow into cancerous (malignant) tumours in the breast ducts or lobules. Most breast cancers are found when they are invasive. This means that the cancer has spread from the breast ducts or lobules into the surrounding breast tissue. Invasive breast cancer can be early, locally advanced or advanced (metastatic). Advanced breast cancer is when cancer cells have spread (metastasised) outside the breast and nearby lymph nodes to other parts of the body. About 5% of breast cancers are advanced when the cancer is first diagnosed.³

Q: How common is breast cancer?

A: About 20,300 people are diagnosed with breast cancer in Australia every year.⁴

Women – In Australia, breast cancer is the most common cancer in women (apart from common skin cancers), with 1 in 7 women diagnosed in their lifetime. Young women can get breast cancer, but it is more common after the age of 40, and the risk increases with age. Pregnant or breastfeeding women can also get breast cancer. See a doctor if you notice any persistent lump during pregnancy.

Men – About 210 men (most aged over 60) are diagnosed with breast cancer each year. It is generally treated in the same way as for women. For more information, visit breastcancerinmen.canceraustralia.gov.au, or bcna.org.au/resource-hub/articles/breast-cancer-in-men.

Transgender, non-binary and gender-diverse – Transgender women taking medicines to boost female hormones have a higher breast cancer risk than men. Trans men and people who have had their breasts removed can still get breast cancer, although the risk is low. For more information, visit bcna.org.au/resources/articles/trans-and-gender-diverse-people-with-breast-cancer.

► See our *LGBTQI+ People and Cancer* booklet.

Q: Does breast cancer run in families?

A: Most people with breast cancer do not have a strong family history, but a small number may have inherited a gene fault (also called a mutation) that increases their breast cancer risk.

BRCA1 and BRCA2 – These are the most common gene mutations linked to breast cancer. Women in families with BRCA1 or BRCA2 are at increased risk of breast and ovarian cancers. Men in families with BRCA2 (and to a lesser extent BRCA1) may be at increased risk of breast, prostate and pancreatic cancers.

Other genes linked to breast cancer – These include ATM, BARD1, CDH1, CHEK2, PALB2, PTEN, RAD51C, RAD51D and TP53. More gene mutations linked to breast cancer are being found all the time. An extended gene panel test checks for the most common ones.

To find out if you have inherited a gene mutation, talk to your doctor about visiting a family cancer clinic or genetic oncologist. Specialists or consulting physicians can also often order genetic tests. Women diagnosed before 50 or with triple negative breast cancer (see page 23) and men with breast cancer should ask for a referral. Genetic testing is covered by Medicare for some people; check with your doctor.

What are the risk factors for breast cancer?


Many factors can increase your risk of breast cancer, but they do not mean that you will develop it. You can also have none of the known risk factors and still get

Personal factors



- Being born female is the biggest risk factor – 99% of breast cancer cases are diagnosed in women.
- Risk increases with age for both men and women.
- More than 3 in 4 breast cancer cases are in women over the age of 50. Free breast screening is available (see page 16).
- Dense breast tissue (as seen on a mammogram) increases your risk.
- Breast implants do not increase breast cancer risk, but some implants are linked with a type of cancer called lymphoma. See tga.gov.au/products/medical-devices/breast-implant-hub.

Lifestyle factors



- Having excess body weight can increase the risk of developing breast cancer after menopause. Keeping your weight within a healthy range can help to reduce this risk.
- The more alcohol you drink, the higher your breast cancer risk. If you choose to drink, the Australian alcohol guidelines suggest you drink no more than 10 standard drinks a week, and no more than 4 standard drinks on any one day.
- Not getting enough exercise or not being physically active.
- Smoking tobacco.

Family history



- About 5–10% of breast cancers are due to an inherited breast cancer gene such as BRCA1 or BRCA2 (see page 9).
- Most people with breast cancer do not have a strong family history. However, having several close relatives (e.g. mother, sister, aunt) on the same side of the family who have had breast or ovarian cancer may increase your risk.
- Having several close relatives on the same side of the family with prostate or pancreatic cancer may increase your risk.

breast cancer. If you are worried, speak to your doctor. For more information, visit canceraustralia.gov.au or www.petermac.org/iprevent.

Hormonal factors




- Taking the oral contraceptive pill for 5 years or more may slightly increase the risk.
- You or your mother taking diethylstilboestrol (DES), a synthetic hormone drug once prescribed during pregnancy.
- Transgender women taking gender-affirming hormones for more than 5 years have a similar risk to those born female.
- Long-term use of menopausal hormone therapy (MHT) containing both oestrogen and progesterone may slightly increase the risk.

Medical history



- Having been previously diagnosed with breast cancer or non-invasive breast conditions (see next page).
- Some non-cancerous conditions of excessive growth of breast cells (atypical ductal hyperplasia or ADH).
- Having radiation therapy to the chest area for Hodgkin lymphoma.
- Males with a rare genetic syndrome called Klinefelter syndrome. Those with this syndrome have 3 sex chromosomes (XXY) instead of the usual 2 (XY).

Reproductive factors



- Never having given birth to a child.
- Starting your first period (menstruating) before the age of 12.
- Being older than age 30 when you gave birth to your first child.
- Never having breastfed a child.
- Going through menopause after the age of 55.

What are the different types of breast conditions and breast cancers?

Non-invasive breast conditions

These are conditions where the abnormal cells have not invaded nearby tissues. Also called carcinoma in situ.

ductal carcinoma in situ (DCIS)	• abnormal cells that look like cells in the breast ducts • may develop into invasive breast cancer • treatment is similar to invasive breast cancer (usually surgery), but chemotherapy is not used
lobular carcinoma in situ (LCIS)	• abnormal cells that look like cells in the breast lobules • classic LCIS does not develop into cancer but it increases the risk of developing it in future • usually treated with surgery, and sometimes hormone therapy • more regular mammograms or other scans needed to check for any changes • other types of LCIS may be treated similarly to DCIS

Invasive breast cancers

Invasive means the cancer cells have grown and spread beyond the breast ducts/lobules and into the surrounding tissue. The 2 main types of invasive breast cancer are named after the cells they resemble.

invasive ductal carcinoma (IDC)	• looks like the cells in the breast ducts • about 80% of breast cancers are IDC
invasive lobular carcinoma (ILC)	• looks like the cells in the breast lobules • about 10% of breast cancers are ILC • can be harder to detect on screening

Rare invasive breast cancers

Inflammatory breast cancer is a rare, fast-growing cancer that affects the lymphatic vessels in the skin of the breast, causing it to become inflamed. Other less common breast cancers include medullary carcinoma, mucinous carcinoma, Paget disease of the nipple (or breast) and papillary carcinoma.

Q: What are the symptoms?

A: Breast cancer sometimes has no symptoms, so regular checks are important (see page 16). Breast changes may not mean cancer, but see a doctor if you notice:

- a lump, lumpiness or thickening, especially in just one breast
- a change in the size or shape of the breast or swelling
- a change to the nipple – change in shape, crusting, sores or ulcers, redness, pain, a clear or bloody discharge, or a nipple that turns in (inverted nipple) when it used to stick out
- a change in the skin – dimpling or indentation, a rash or itchiness, scaly appearance, unusual redness or other colour changes
- swelling or discomfort in the armpit or near the collarbone
- ongoing, unusual breast pain not related to your period.

Q: Which health professionals will I see?

A: You may be sent for tests after a screening mammogram (see page 16), or your general practitioner (GP) may arrange tests to check your symptoms. If these tests do not rule out cancer, you will usually be referred to a specialist or breast clinic. If breast cancer is diagnosed, you will see a breast surgeon or a medical oncologist, who will talk to you about your treatment options. Often these will be discussed with other health professionals at a multidisciplinary team (MDT) meeting. During and after treatment, you will see a range of health professionals who specialise in different aspects of your care (see pages 14–15). You may not see all members of the MDT.



For an overview of what to expect at every stage of your cancer care, visit cancer.org.au/cancercareguides/breast-cancer. This is a short guide to what is recommended, from diagnosis to treatment and beyond.

Health professionals you may see

GP	arranges initial tests to investigate symptoms; assists you with treatment decisions and works in partnership with your specialists in providing ongoing care when cancer is diagnosed
breast physician	diagnoses breast cancer and other breast conditions, especially when initial test results are unclear; provides care and support during and after treatment
radiologist	analyses mammograms, ultrasounds and other scans; performs fine needle and core biopsies to confirm diagnosis
radiographer/ sonographer	performs mammograms, breast ultrasounds and other scans
breast surgeon	diagnoses breast cancer, performs surgical biopsies in some clinics; performs breast surgery; some breast surgeons also perform breast reconstruction; some breast surgeons specialise in using plastic surgery techniques to reconstruct breast tissue after surgery, also called oncoplastic surgery
reconstructive (plastic) surgeon	performs breast reconstruction after mastectomy
radiation oncologist	treats cancer by prescribing and overseeing a course of radiation therapy
medical oncologist	treats cancer with drug therapies such as chemotherapy, hormone therapy, targeted therapy and immunotherapy (systemic treatment)
breast care nurse	provides breast cancer care; also provides information and facilitates referrals during and after treatment
chemotherapy nurse	administers drugs and provides care, information and support throughout treatment

anaesthetist	assesses your health before the operation, administers anaesthetic and looks after you during and after surgery; plans your pain relief
radiation therapist	plans and delivers radiation therapy
physiotherapist, exercise physiologist	help restore movement and mobility, and improve fitness and wellbeing
occupational therapist	assists in adapting your living and working environment to help you resume usual activities after treatment
lymphoedema practitioner	educates people about lymphoedema prevention and management, and provides treatment if lymphoedema occurs; is often a physiotherapist or occupational therapist
social worker	links you to support services and helps you with emotional, practical and financial issues
dietitian	helps with nutrition concerns and recommends changes to diet during treatment and recovery
psychologist, counsellor	help you manage your emotional response to diagnosis and treatment
genetic counsellor	provides advice for people with a strong family history of breast cancer or for people with a genetic condition linked to cancer
sexual health counsellor	helps you manage the effects of cancer and its treatments on your sex life

Diagnosis

If you notice breast changes or swelling in your armpit, your GP will ask about your medical history and any family history of breast cancer. They will do a physical examination, checking both breasts and the lymph nodes in your armpit and above your collarbone. They may also arrange imaging tests and a biopsy. This is called a triple test. Sometimes a specialist will arrange extra tests.

If you feel uncomfortable having a test, you can ask for a nurse, health professional, friend or family member to be in the room with you or, if that is not possible, to be nearby.

Mammogram

A mammogram is a low-dose x-ray of the breast tissue. It can show changes that are small or cannot be felt during a physical examination. Let staff know before the mammogram if you have breast implants.

Your breast is placed between 2 x-ray plates. The plates press together firmly to spread out the breast tissue so that clear pictures can be taken. You will feel some pressure, which can be uncomfortable, but the mammogram only takes 10–15 seconds. Both breasts will be checked.

A national screening program provides a free mammogram every 2 years for:

- all women over 50
- trans women over 50 who have been taking gender-affirming hormones for 5 years or more
- trans men over 50 who have not had gender-affirming chest surgery.



Before a scan, tell the doctor if you have any allergies or if you have had a reaction to dyes during previous scans. You should also tell them if you have diabetes or kidney disease or you are pregnant or breastfeeding.

Those over 40 can also book a free mammogram. Speak to your doctor about whether breast screening before 50 is right for you. For more information, call BreastScreen Australia on 13 20 50 or visit health.gov.au/bsa.

Tomosynthesis – Also known as three-dimensional mammography, tomosynthesis takes x-rays of the breast from many angles and combines them into a three-dimensional (3D) image. This may be better for finding small breast cancers, particularly in dense breast tissue.

Contrast enhanced mammogram (CEM) – This combines tomosynthesis with a dye (contrast) that is injected into a vein in your arm. A CEM may be helpful for people with dense breast tissue.

Ultrasound

An ultrasound uses soundwaves to create a picture of breast tissue. It's often the first test done in women under 30 with breast changes, or if a routine screening mammogram has found breast changes, or you or your GP feel a lump.

A gel will be spread on your breast, and then a small device (transducer) is moved over the breast and armpit. This sends soundwaves that echo when they meet something dense, like a tumour. A computer creates a picture from these echoes. The scan takes 15-20 minutes and is painless.

Breast MRI scan

A magnetic resonance imaging (MRI) scan uses a large magnet and radio waves to take pictures of the breast tissue. It does not use radiation. It is mainly used for people at high risk of breast cancer or who have very dense breast tissue or breast implants. It may also be used if other imaging test results are unclear or to help plan surgery.

Before a breast MRI scan, you will usually have an injection of a dye (contrast) to help show any abnormal breast tissue. You will lie face down on a table, which will slide into a large, cylinder-shaped machine. The scan can take up to 40 minutes. It is painless but loud, so you will wear headphones. Some people feel claustrophobic. If you are concerned, talk to your doctor beforehand. You may be offered a mild sedative to take before the scan.

Biopsy

In a biopsy, a small sample of cells or tissue is taken from the lump or area of concern. A specialist doctor called a pathologist then checks the sample under a microscope to see if there are any cancer cells.

During the biopsy, the doctor might place a small metallic clip to mark the area. You should let the medical team know if you take blood-thinning medicine as you will need to stop it 5-7 days before the procedure. Your breast may be bruised for a few days afterwards.

There are different ways of taking a biopsy and you may need more than one type. The biopsy may be done in a specialist's rooms, at a radiology practice, in hospital or at a breast clinic.

- **Fine needle aspiration (FNA)** – A thin needle is inserted into an abnormal lymph node or tissue, often with an ultrasound to guide

it into place. Tiny pieces of tissue can then be sucked out through the needle. A local anaesthetic may be used to numb the area.

- **Core biopsy** – Several pieces of tissue are removed with a needle. Local anaesthetic numbs the area, and you may have a mammogram, ultrasound or MRI scan to guide the needle into the right area.
- **Vacuum-assisted core biopsy** – A needle attached to a suction-type instrument is inserted into the breast through a small cut in the skin. A larger amount of tissue is removed with a vacuum biopsy, making it more accurate in some cases. The needle is usually guided into place with an ultrasound or MRI. A local anaesthetic is used, but you may feel some discomfort. Stitches are not usually needed.
- **Surgical (excision) biopsy** – If a needle biopsy is not possible, or the diagnosis is unclear, you may have a surgical biopsy to remove all or part of a lump. A wire or small surgical clip may be inserted to act as a guide during the surgery. The tissue is then removed under general anaesthetic. This is usually done as day surgery.

Further tests

If these tests show that you have breast cancer, you may have further tests to check whether the cancer has spread to other parts of your body. You will have a blood test to check your general health and, in some cases, it will test for specific tumour markers.

You may also have some of the following types of scans.

Bone scan – A bone scan is used to see if the breast cancer has spread to your bones. A small amount of radioactive solution is injected into a vein, usually in your arm. This solution is attracted to abnormal areas of the bone. After a few hours, the doctors look at the bones with a scanning machine. The scan is painless and the solution is not harmful.

CT scan – A CT (computerised tomography) scan uses x-ray beams to take pictures of the inside of the body. Before the scan, dye (contrast) will be injected into a vein in your arm. This helps to make the pictures clearer. You might have a feeling of passing urine, but this won't last long. You then lie flat on a table while the scanner takes pictures. The scan takes about 30 minutes and is painless.

PET scan – In a PET (positron emission tomography) scan, a small amount of low-level radioactive solution is injected into a vein in the arm or hand. Any cancerous areas take up more of the radioactive solution and may show up brighter in the scan. You will usually have to fast (not eat anything) for 4–6 hours before the scan.

Staging breast cancer

These tests show the size of the breast cancer and whether it has spread to other parts of the body. This is called staging. It helps you and your healthcare team decide what treatment will be best.

The most common staging system used for breast cancer is the TNM system. Letters and numbers describe how big the tumour is (T), if cancer has spread to nearby lymph nodes (N), or if it has spread to the bones or other organs, which is known as having metastasised (M).

The staging system also describes details about the tumour such as oestrogen and progesterone receptor status, HER2 status (see pages 22–23) and the grade (see opposite page). Staging may be done before or after surgery, depending on when the doctors have full information about the cancer. The cancer may be classified as:

- **Early breast cancer (stage 1 or 2)** – The cancer is contained in the breast and may or may not have spread to lymph nodes in the armpit.

Grading breast cancer

The grade describes how active the cancer cells are and how fast the cancer is likely to be growing.

grade 1 (low grade)	Cancer cells look a little different from normal cells. They are usually growing slowly.
grade 2 (intermediate grade)	Cancer cells do not look like normal cells. They are growing faster than grade 1 cancer cells.
grade 3 (high grade)	Cancer cells look very different from normal cells. They are usually growing fast.

- **Locally advanced breast cancer (stage 3)** – The cancer is larger than 5 cm, has spread to tissues around the breast such as the skin or muscle or ribs, or has spread to a large number of lymph nodes.
- **Metastatic breast cancer (stage 4)** – The cancer has spread to other parts of the body from the breast. Also called secondary or advanced breast cancer, it's different from locally advanced breast cancer.

Prognosis

Prognosis means the expected outcome of a disease. You can talk to your doctor about what might happen, but it's not possible for anyone to predict the exact course of the disease. To work out your prognosis, your doctor will consider the stage and grade of the cancer, as well as features such as the hormone receptor and HER2 status (see pages 22-23).

The survival rates for people with breast cancer have increased significantly due to more people taking part in screening, better tests and scans, and improved treatments. Compared to other cancers, breast cancer has one of the highest survival rates when diagnosed early.

Tests on breast tissue

If tests on the biopsy sample confirm you have breast cancer, extra tests on the biopsy sample will be done to understand more about the breast cancer and help plan treatment. The results will be included in the pathology report.

Hormone receptor status ER+ and/or PR+



70–80%
of all breast
cancers

The hormones oestrogen and progesterone are produced naturally in the body. A receptor is a protein on the surface of the cell. Normal breast cells have oestrogen receptors (ER) and progesterone receptors (PR). Breast cancers that have these receptors are known as ER positive (ER+) and/or PR positive (PR+). Oestrogen may stimulate the cancer cells to grow. Progesterone receptors are important in normal breast cells but are not usually used to guide cancer treatment.

ER+ cancers are usually treated with hormone therapy drugs (also known as endocrine therapy) that block the receptor, or drugs that reduce the amount of hormones that the body makes (aromatase inhibitors).

If the cancer has low levels of oestrogen receptors, hormone therapy drugs will still be used. These drugs are not used for cancers with no oestrogen receptors.

HER2 status HER2+



15–20%
of all breast
cancers

HER2 (human epidermal growth factor receptor 2) is a protein that is found on the surface of some cells and controls the way cells grow and divide.

Tumours with high levels of these receptors are called HER2 positive (HER2+). Tumours with low levels are called HER2 negative (HER2– or HER2 low).

It's usually recommended that people with HER2+ breast cancer have chemotherapy and targeted therapy before they have surgery (neoadjuvant treatment, see page 28). Depending on how the cancer responds to the neoadjuvant treatment and surgery, you may also have chemotherapy or targeted therapy after surgery (adjuvant treatment, see page 28).

Triple negative breast cancer ER–, PR– and HER2–



10–20%
of all breast
cancers

Some breast cancers do not have oestrogen (ER–), progesterone (PR–) or HER2 (HER2–) receptors. These are called triple negative breast cancers.

Triple negative breast cancers do not respond to hormone therapy or to the targeted therapy drugs used for HER2+ cancers in early breast cancers.

These types of cancer usually respond well to chemotherapy, so this may be used before and/or after surgery (neoadjuvant/ adjuvant treatment, see page 28).

Some other types of targeted therapy may be used for triple negative breast cancer.

Recently, various types of immunotherapy have been shown to work well for some triple negative cancers. These may be used before surgery for larger cancers or for cancers that also affect lymph nodes.

What are gene expression profile tests?

Gene expression profile tests may be done on the biopsy sample. These tests look at which genes are active in the cancer cells. The results give information about the risk of the cancer coming back.

These tests may also be called genomic tests or molecular assays. They are different to genetic tests, which are used to look for inherited gene faults (see page 9).

The gene expression profile tests available in Australia are Oncotype DX, EndoPredict, PAM50, Prosigna and MammaPrint.

The test results will help the doctor work out if chemotherapy will be helpful after surgery. It can take 14 days for the results to come back, so if you need these tests it may be helpful for your surgeon to order them before you see your oncologist. They would usually do this after discussing it in an MDT meeting (see page 13).

Ask your doctor if a gene expression profile test is an option for you. The standard pathology tests done on all breast cancers may be all that is needed for your treatment plan.

Gene expression profile tests may not be covered by Medicare; check what you may have to pay.

Key points about diagnosing breast cancer

Main tests

Tests to diagnose breast cancer include:

- physical examination
- mammogram (breast x-ray)
- ultrasound
- breast MRI scan
- biopsy (taking a tissue sample).

Further tests

Further tests can give more information about the cancer to help guide treatment. These tests may include:

- blood tests
- bone scan
- CT scan
- PET scan.

Staging

The stage shows how far the cancer has spread. Early breast cancer is stage 1 or 2. Locally advanced breast cancer is stage 3. Cancer that has spread to other parts of the body is stage 4.

Key information about the cancer

- Hormone receptor status (ER+/- and PR+/-) shows whether the cancer may respond to hormone therapy. ER status is particularly important for planning treatment.
 - HER2 status (HER2+/-) shows whether the cancer may respond to a combination of chemotherapy and targeted therapy or chemotherapy and hormone therapy.
 - Gene expression profile tests (also called genomic tests or molecular assays) look at gene patterns in cancer cells. They can help show if chemotherapy would be useful in early breast cancer but they are not always needed.
-



Making treatment decisions

Sometimes it is difficult to decide on the type of treatment to have. You may feel that everything is happening too fast, or you might be anxious to get started.

Check with your specialist how soon treatment should begin, as it may not affect the success of the treatment to wait a while. Ask them to explain the options, and take what time you can before making a decision.

Know your options – Understanding the disease, the available treatments, possible side effects and any extra costs can help you weigh up the options and make a well-informed decision. Check if the specialist is part of a multidisciplinary team (see page 13) and if the treatment centre is the most appropriate one for you – you may be able to have treatment closer to home, or it might be worth travelling to a centre that specialises in a particular treatment.

Record the details – When your doctor first says you have cancer, you may not remember everything you are told. Taking notes can help. If you would like to record the discussion, ask your doctor first. It's also a good idea to have a family member or friend go with you to appointments to join in the discussion, write notes or simply listen.

Ask questions – If you are confused or want to check anything, it's important to ask your specialist questions. Try to prepare a list before appointments (see page 67 for suggestions). If you have a lot of questions, you could talk to a breast care nurse.

Consider a second opinion – You may want to get a second opinion from another specialist, to confirm or clarify your specialist's recommendations, or to reassure you that you have explored all of your options. Specialists are used to people doing this. Your GP or specialist can refer you to another specialist and send your initial results to that person. You can get a second opinion even if you have started treatment or still want to be treated by your first doctor. You might decide you would prefer to be treated by the second specialist.

It's your decision – Adults have the right to accept or refuse any treatment that they are offered. For example, some people with advanced cancer choose treatment that has significant side effects, even if it gives only a small benefit for a short period of time. Others decide to focus their treatment on quality of life. You may want to discuss your decision with the treatment team, or your GP, family and friends.

► See our *Cancer Care and Your Rights* fact sheet.

Should I join a clinical trial?

Your doctor or nurse may suggest you take part in a clinical trial. Doctors run clinical trials to test new or modified treatments and ways of diagnosing disease to see if they are better than current methods. For example, if you join a randomised trial for a new treatment, you will be chosen at random to receive either the best existing treatment or the modified new treatment. Over the years, trials have improved treatments and

led to better outcomes for people diagnosed with cancer.

You may find it helpful to talk to your specialist, clinical trials nurse or GP, or to get a second opinion. If you decide to take part in a clinical trial, you can withdraw at any time. For more information, visit australiancancertrials.gov.au.

► See our *Understanding Clinical Trials and Research* booklet.

Treatment

This chapter focuses on treatment for early and locally advanced breast cancer. For information on the treatment of advanced breast cancer, see page 50.

Treatment for early or locally advanced breast cancer varies. The treatment that's best for you will depend on your test results, where the cancer is in the breast, the stage and grade of the cancer (see pages 20–21), and whether the cancer is hormone receptor and/or HER2 positive or triple negative (see pages 22–23). Your treatment team will also consider your age and general health, and your preferences.

Treatment for early and locally advanced breast cancer usually includes surgery. Before surgery, you may have other types of treatment to shrink the cancer. This is called neoadjuvant treatment (see next page).

Surgery

The type of surgery your doctor suggests will depend on the type and stage of the cancer, where it is in the breast, the size of your breast, and what you prefer. In most cases, you will have one or more lymph nodes removed from the armpit (called axillary surgery, see page 32). Some people also choose to have a new breast shape made during the operation (breast reconstruction, see page 31).

The 2 different types of surgery used for breast cancer are:

- **breast-conserving surgery** – when only part of the breast is removed (see page 29)
- **mastectomy** – when the whole breast is removed (see page 30).

Depending on your situation, you may have a choice between the 2 types of surgery. Research has shown that for most early breast cancers, having breast-conserving surgery followed by radiation therapy (see pages 40–42) works just as well as a mastectomy.

The operations have different benefits, risks and side effects. Talk to your doctor about the best option for you.

Treatments and surgery

Treatment before surgery

While surgery is often the main treatment for both early and locally advanced breast cancer, you may have other treatments before surgery.

This is called neoadjuvant treatment and it may be discussed by an MDT (see page 13). Chemotherapy is often used before surgery or you may have targeted therapy or immunotherapy (see pages 47–49), or a combination of these treatments. Occasionally you might also have hormone therapy (see page 45–47) before surgery.

Treatment can help to reduce the size of the cancer before surgery and improve your chance of a good outcome. It may also mean you can have less complex surgery or you can choose to have breast-conserving surgery instead of a mastectomy.

In some cases – particularly for people with HER2+ or triple negative cancers – neoadjuvant treatment can kill all cancer cells. Called a complete pathological response, it improves the chance of a good outcome.

Ask your doctor if neoadjuvant treatment is an option for you. People with early breast cancer may find the Neoadjuvant Patient Decision Aid helpful. For more information, visit myneoguide.com.

Treatment after surgery

After surgery, you may have radiation therapy, chemotherapy, hormone therapy, targeted therapy, immunotherapy or a combination of all these treatments. This is called adjuvant treatment. It helps to destroy any tiny cancer cells that may remain after surgery.

Breast-conserving surgery

Removing only part of the breast is called breast-conserving surgery. It's also known as a lumpectomy or wide local excision.

The surgeon removes the tumour and some of the healthy tissue around it, so that you can keep as much of your breast as possible. The operation will leave a scar, and it may change the size and shape of the breast and the position of the nipple.

Pathology tests on breast tissue

A pathologist looks at the removed tissue under a microscope to check for an area of healthy cells around the cancer (called a clear margin).

The pathologist will also give information about:

- the size and grade of the cancer (see pages 20–21)
- whether the cells are hormone receptor positive and/or HER2+ or triple negative (see pages 22–23)
- whether the cancer has spread to any lymph nodes (see pages 32–33).

If your removed tissue shows multiple cancers, each cancer will be tested separately. The pathology report will help your doctors work out what other treatment may be best for you. If there are cancer cells found close to the edge of the tissue (which is called an involved or positive margin), there is a higher risk of the cancer returning.

You may need to have further surgery to remove more tissue (called a re-excision or wider excision). Your doctor may also suggest that you have a mastectomy (see page 30).

After having breast-conserving surgery, even if you have a clear margin, you will usually have radiation therapy to destroy any cancer cells that may be left in the breast or armpit (see pages 40–42).

Mastectomy

Surgery to remove the whole breast is called a mastectomy. One breast may be removed (single or unilateral mastectomy) or both breasts (double or bilateral mastectomy). A mastectomy may be recommended if:

- there is cancer in more than one area of the breast
- the cancer is large compared with the size of the breast
- it is difficult to get a clear margin around the tumour
- you have inflammatory breast cancer (see page 12)
- you have had radiation therapy to that breast and cannot have it again
- you have the BRCA1 or BRCA2 or another breast cancer gene mutation.

If the cancer has come back, or there is a new cancer in the same breast, you may still be able to keep your breast. Or you may choose to have a mastectomy, even if you have a small cancer. Talk to your doctor. You don't usually need radiation therapy after a mastectomy.

Having a mastectomy can be confronting but there are several options for your breast and chest appearance (see page 37). In some cases, the surgeon may be able to do a skin-sparing or nipple-sparing mastectomy, where more of the skin (with or without the nipple) is kept. Talk to your surgeon about whether this is an option. If you have a reconstruction (see opposite page), and are able to have a skin-sparing or nipple-sparing mastectomy, they can sometimes be done at the same time.

If you decide not to have a reconstruction, you will have a flat closure. You can also have an immaculate or aesthetic flat closure, a procedure where the skin is smoothed out. If you want to, you can then also wear a soft breast shape (form) in a special bra while you heal. Breast Cancer Network Australia (BCNA) provides a free bra and soft form. Speak to your breast care nurse for details. After you have healed, if you want to, you can be fitted for a permanent breast prosthesis (see page 55).

What about the other breast?

If you need a mastectomy because of cancer in one breast, you may think it's safer to have the other breast removed as well. For most people, the risk of getting cancer in the other breast is low.

If you have the BRCA1 or BRCA2 gene, or another rare breast cancer gene mutation, this does increase

the risk of developing another breast cancer, so you may choose to have a double mastectomy (bilateral mastectomy) to remove both breasts.

Deciding whether to have a double mastectomy is a complex decision. It's best to talk with your treatment team about the risks and benefits before making a final decision.

Breast reconstruction

Breast reconstruction is surgery to make a new breast shape (also called a breast mound). There are different ways to construct a breast shape.

It can be done using:

- a breast implant
- a flap of your own skin or fat (an autologous reconstruction)
- a breast implant and your own tissue.

A breast reconstruction can sometimes be done at the same time as a mastectomy (immediate reconstruction); or you may need or choose to wait for several months or years before having a reconstruction (delayed reconstruction). If you are not having an immediate reconstruction but might consider it in the future, discuss this with your surgeon before surgery. This will help them to plan the mastectomy.

Some people decide not to have a reconstruction and/or wear a breast prosthesis (see page 55).

► See our resources on breast prostheses and reconstruction.

Removing lymph nodes (axillary surgery)

Cancer cells that spread from the breast usually spread first to the axillary lymph nodes, which are in and around the armpit. Removing some or all of these lymph nodes helps your doctor to check if any cancer has spread. The operation to remove lymph nodes is called axillary surgery. It's usually done during breast surgery but may be done in a separate operation. There are 2 main types of axillary surgery.

Sentinel lymph node biopsy (SLNB) – When breast cancer spreads outside the breast, it first goes to a particular lymph node or nodes in the armpit or near the breastbone (sternum). These are called the sentinel nodes. A sentinel node biopsy removes these nodes so they can be tested for cancer cells (see opposite page).

If there are no cancer cells in the sentinel nodes, no more lymph nodes are removed. If there is more than a small amount of disease in the sentinel nodes, you may have axillary lymph node dissection (see below) or radiation therapy (see pages 40–42).


Axillary lymph node dissection (ALND) – If cancer is found in multiple lymph nodes, then many or all of the axillary lymph nodes (usually 10–25) may be removed to reduce the risk of the cancer coming back (recurrence). The nodes are tested and the results guide what other treatment may be needed. ALND is also called axillary lymph node clearance (AC). If the cancer is only in a few lymph nodes, you will probably have radiation therapy instead.

Side effects – You may have numb, stiff, weak, tingling or painful arms or shoulders; a numb armpit and chest; fluid collecting near the scar (seroma); lymphoedema (see page 52); and cording (see page 53). Side effects are worse after ALND because more lymph nodes are removed.

Finding the sentinel nodes

To work out which lymph nodes are sentinel nodes, one or a combination of these procedures is used:

1. Lymphatic mapping




A small amount of a harmless radioactive solution is injected into the skin over the breast cancer tumour.

A CT scan is then taken to show which lymph nodes the radioactive solution flows to first. These are most likely to be the sentinel nodes.

Lymphatic mapping is done either the day before or on the day of the surgery.


2. Dye injection (not always used)



If dye is being used, it will be injected into the breast. The dye, which may be blue or green, moves into the lymphatic vessels and stains the sentinel nodes first. This is done under general anaesthetic during the surgery.

Because of the dye, you may notice blue-green urine (wee) and bowel movements (poo) when you go to the toilet the next day. You may also have a blue patch on the breast for weeks or longer. Your skin may look a bit grey but will fade once the dye washes out in your urine.

3. Handheld probe




As well as looking at where the dye travels to first (if it's used), the surgeon uses a small handheld device called a probe during the surgery to detect the radioactive solution injected during the lymphatic mapping.

This helps to check that the sentinel nodes have been located and the surgeon can then remove them for testing.

What to expect after surgery

If you have any questions about your recovery and how best to look after yourself when you get home, ask the doctors and nurses caring for you. Ask for a referral to a breast care nurse; they can give you information about what to expect after surgery and provide support.

Managing dressings and tubes



A dressing will cover the wound to keep it clean. This may be changed while you are in hospital but is usually removed after about a week. You may have one or more drainage tubes to drain fluid from the surgical site into a bottle or bag. These can stay in place for up to one week, or occasionally 2 weeks. Nurses will show you how to look after the wound and drainage tubes at home, or a community nurse, GP or your surgeon may help you. If you notice redness or discharge around the surgical area or develop a fever over 38°C, let your treatment team know immediately.

Recovery time



The time it takes to recover from surgery will depend on the type of surgery you have had and your health. You may feel better after a few days, or it may take several weeks or longer if you have had a mastectomy with a reconstruction.

Avoid heavy lifting



Do not do vigorous physical activity or heavy lifting in the initial weeks after surgery. Your treatment team will let you know when you can go back to normal activities. You may be given some gentle exercises to reduce the risk of shoulder stiffness.

Shower carefully



Keep the wound clean, and gently pat it dry after showering. Avoid bathing and swimming for at least a week.

Your hospital stay will depend on the surgery you have and how well you recover:

- **breast-conserving surgery** – you usually go home the same day, or may stay overnight
- **mastectomy** – you usually stay in hospital for 1–2 nights
- **reconstruction after mastectomy** – you usually stay in hospital for several days.

Manage pain



While in hospital, you will have pain relief through a drip (intravenous or IV), an injection or as tablets. You will also be given pain medicine when you go home. You are likely to need stronger pain relief after a lymph node dissection (see page 32) or a mastectomy than after breast-conserving surgery.

Prevent blood clots



Apply moisturiser



Gently massage the area with moisturiser once any stitches or adhesive strips are removed and the wound has completely healed. About 6 weeks after surgery, your surgeon may suggest that you use silicone gels and sheets to reduce scarring.

Avoid cuts



Your treatment team may advise you to wait until the wound has completely healed if you want to shave or wax your armpits.

While in bed, you should try to do some deep breathing exercises, and move your legs around to help prevent blood clots in the deep veins of your legs (deep vein thrombosis or DVT). As soon as you are able, you will be asked to get up and walk around. You may wear elastic (compression) stockings or use other devices to help prevent clots. Your doctor might prescribe medicine that reduces the risk of blood clots forming.

What your breast looks like after surgery

How your breast will look after surgery depends on the type of surgery you have, the size of your breast and your body shape. It can take a few weeks for any bruising and swelling of the surgery area to go away.

After breast-conserving surgery – The size and position of the scar will depend on how much tissue is removed. The scar will usually be near where the cancer was, around the areola or near the breast fold. But this varies depending on your breast size and how much tissue needs to be removed. If a larger area needs to be removed, techniques known as oncoplastic surgery can reshape the breast after the surgery.

After a mastectomy – The scar position will depend on the type of mastectomy you have. After a skin-sparing mastectomy, the scar will be where your nipple was; it may be horizontal or vertical. If you have flat closure, it will run across the centre of your chest. After a nipple-sparing mastectomy, it will run in the fold under the base of the breast or around the nipple. If you have lymph nodes removed, there may also be a scar in the armpit. At first the scar will be firm, slightly raised and red. Over the next few months, it will flatten and fade.

Impact on self-esteem

Scars or changes to how your breast looks can affect how you feel about yourself (self-image and self-esteem, see page 56). If you have had a mastectomy or part of your breast removed, it's common to feel a sense of loss. It may also affect your sense of identity.

Seeking support – It can help to talk to someone who has had breast cancer surgery. Cancer Council's Peer Connect program may be able to link you to others with a similar experience. Talking to a counsellor or psychologist can also help. Call Cancer Connect on 13 11 20.

Breast appearance after surgery

What your breast will look like after surgery depends on the type of surgery you have, how much tissue is removed and your body shape. These pictures give some examples, but ask your surgeon for photographs to help you choose the best approach for you. Some procedures may not be suitable for you.



Breast-conserving surgery

Two years after surgery to the left breast. The scar can still be seen but has healed and faded a little. The size of the scar varies but can be around 10 cm. The surgery was followed by radiation therapy.



Mastectomy with flat closure

The left breast has been removed and the chest closed in an aesthetic flat closure, a procedure where the excess skin and tissue are removed and the chest is smoothed out. The scar runs across the chest wall. This photo is some time after surgery, once the scar has healed.



Nipple-sparing mastectomy with an implant reconstruction

The left breast has been removed with a nipple-sparing mastectomy, followed by a breast reconstruction using an implant. If it is not possible to keep the nipple, there is the option of having a nipple reconstruction later.



Double mastectomy with a flap reconstruction

Both breasts have been removed in a skin-sparing mastectomy. This was followed by a DIEP flap breast reconstruction, using skin and tissue from the stomach, and then later a nipple reconstruction and tattoo.

Side effects of surgery

Some common side effects are discussed below. Talk to your treatment team about the best ways to deal with them. For more information, see the *Managing side effects* chapter (pages 52–57).

Fatigue – Cancer treatment and the emotional impact of the diagnosis can be tiring. Fatigue is common and may continue for weeks or months. Research shows that exercise during and after cancer treatment is safe and can help improve fatigue. Cancer Connect may be able to connect you with an exercise program near you – call 13 11 20. You can also ask your GP if you are eligible for some Medicare-funded sessions with an exercise physiologist or a physiotherapist.

- ▶ See our *Understanding Fatigue and Cancer* fact sheet and *Exercise for People Living with Cancer* booklet.

Shoulder stiffness – Arm and shoulder pain, weakness, stiffness and reduced movement are common after surgery and after radiation therapy. Ask your treatment team when you can start exercising your arm. A physiotherapist or exercise physiologist can show you exercises to reduce shoulder stiffness or pain. This may help prevent lymphoedema (see page 52).

- ▶ See our *Arm and Shoulder Exercises* fact sheet.

Numbness and tingling – Surgery can bruise or injure nerves. You may feel numbness and tingling in the armpit, upper arm or chest area. You



McGrath Breast Care Nurses are registered nurses specifically trained to support people with breast cancer from the time of diagnosis and throughout treatment. There are more than 400 McGrath nurses across Australia. For more information, visit mcgrathfoundation.com.au.

may also notice a loss of feeling in your breast or nipple. These changes often improve within a few weeks but may take longer. Sometimes the numbness or tingling may not go away completely. A physiotherapist or occupational therapist can give you exercises that may help.

Seroma – Fluid may collect in or around the surgical scar and cause a balloon-like swelling. This is most common after a mastectomy. A seroma can also develop in the armpit after an ALND (see page 32). The build-up of fluid can be uncomfortable but is not harmful. Some breast care nurses, your specialist or GP, or a radiologist can drain the fluid using a fine needle and a syringe. This procedure is not painful, but it may need to be repeated over a few appointments.

Lymphoedema – Fluid building up in the tissue of the arm or breast may cause swelling after any lymph node surgery. It's common to have some swelling of your arm or breast after surgery, but this usually settles in the weeks afterwards. If this swelling builds up over weeks or months, this usually means you have lymphoedema. It can happen any time, even years after surgery (or radiation therapy) to the lymph nodes. See page 52 for ways to prevent and manage the symptoms of lymphoedema.

► See our *Understanding Lymphoedema* fact sheet.

Post-mastectomy pain – It's rare to have prolonged pain after a mastectomy but you may find the scar uncomfortable for some time. If pain or discomfort is ongoing, let your treatment team know.

Cording – Also known as axillary web syndrome, cording is caused by hardened lymph vessels. It feels like a tight cord running from your armpit down the inner arm, sometimes to the palm of your hand. See page 53 for ways to manage it.

Radiation therapy

Also known as radiotherapy, radiation therapy uses a controlled dose of radiation to kill cancer cells or damage them so they cannot grow, multiply or spread. The radiation is usually in the form of focused, high-energy x-ray beams. It does not cause you to become radioactive during the period of treatment.

Radiation therapy may be recommended:

- after breast-conserving surgery – it's usually part of standard treatment
- after a mastectomy – you may have radiation to the chest wall and lymph nodes above the collarbone, and sometimes to the lymph nodes next to the breastbone
- if the sentinel node has cancer cells – you may have radiation to the armpit instead of ALND (see page 32)
- after neoadjuvant chemotherapy and before surgery (see pages 27–31)
- after adjuvant chemotherapy (see pages 43–44).

You will usually start radiation therapy within 8 weeks of surgery. If you're having chemotherapy after surgery, radiation therapy will begin about 4 weeks after chemotherapy has finished.

Planning radiation therapy

Treatment is carefully planned to cause the most harm to the cancer cells and to limit damage to the surrounding healthy tissues. Planning involves several steps, which may be over a few visits.

You will have a planning session at the radiation therapy centre. During this appointment, you will have a planning CT scan of the area to be treated. Sometimes marks are put on your skin so the radiation therapists can make sure you are lined up correctly each time you are treated. These marks are usually tiny permanent dots (tattoos), about the size

of a freckle. If you have had breast-conserving surgery, the surgeon can sometimes place tiny markers (called fiducial markers) in your breast tissue to show where the cancer used to be. This helps the radiation oncology team to deliver the radiation therapy more precisely.

You may be asked to try a deep inspiration breath hold (DIBH) technique. This involves taking and holding a deep breath for 20–30 seconds during treatment. The technique helps to inflate the lungs and move the heart away from the radiation field, reducing the risk of heart damage.

Having radiation therapy

You will probably have radiation therapy daily from Monday to Friday for 1–4 weeks. Most people have radiation therapy as an outpatient and go to the treatment centre each day.

Each radiation therapy session will be in a treatment room. Setting up the machine can take 10–30 minutes, but the actual treatment takes only 1–5 minutes. You will lie on a table under the machine and your breast will be exposed. The radiation therapist will leave the room and then switch on the machine, but you can talk to them through an intercom.

Radiation therapy is not painful, but you will need to lie still while it is given. Most people will be lying on their back with their arms up. If the DIBH technique is recommended for you, the radiation beam will only be turned on once you have taken the deep breath.

If you are having radiation therapy at a private centre, Medicare will cover some of the cost, but your private health insurance may not, so you may have to pay some of the cost yourself (out-of-pocket costs). If you are worried about the cost, speak to your treatment team about having treatment in a public hospital.

Side effects of radiation therapy

Radiation therapy may cause the following side effects:

Skin problems – You may have some redness around the treated area. The skin may become dry and itchy, blister, or become moist and weepy. It usually returns to normal 4–6 weeks after radiation therapy ends. Sometimes skin can become very irritated or peel (radiation dermatitis). You may need dressings, or special creams or gels, to help the area heal.

Tiredness – You may start to feel tired 1–2 weeks after radiation therapy begins. Fatigue usually gets better a few weeks after treatment finishes.

Aches – You may feel minor aches or shooting pain in the breast area during or after radiation therapy. It should ease over time.

Swelling – Some people have swelling or fluid build-up in the breast (breast oedema or lymphoedema) that can last for up to a year or longer. Radiation therapy to the armpit increases the risk of swelling in the arm (lymphoedema, see page 52).

Hair loss – Radiation therapy to the breast won't make you lose the hair on your head, but you will usually lose hair from the treated armpit.

Other side effects – Occasionally, late effects can develop months or years after radiation therapy has finished. The most common late effect is fibrosis, or scar tissue, inside the breast, which can make it feel firmer or tender. There is also a slight risk of lung and heart problems, particularly if you have other risk factors, such as if you smoke cigarettes. In rare cases, radiation therapy may cause a second cancer years later.

► See our *Understanding Radiation Therapy* booklet.

Chemotherapy

Chemotherapy uses drugs to kill cancer cells or slow their growth. It may be used before or after surgery (see page 28). It's often used for breast cancers that are not sensitive to hormone therapy, are HER2+ or triple negative, or for inflammatory breast cancers (see page 12). Chemotherapy is also sometimes used for hormone receptor positive breast cancers.

Having chemotherapy

There are different types of chemotherapy drugs. The choice of drugs will depend on the type of cancer, how far it has spread and what other treatments you are having. Your medical oncologist will talk to you about the most suitable types of chemotherapy drugs, as well as their risks and side effects. For more information, visit eviq.org.au.

Chemotherapy for breast cancer is usually given through a vein (intravenously) or through a port. This is a thin plastic tube that is surgically inserted under the skin, usually in the chest or arm, under local anaesthetic. It stays in your vein throughout the treatment course.

You will usually be treated as an outpatient, but occasionally you may have to stay in hospital overnight. Chemotherapy is usually given once every 1-3 weeks for 3-6 months.

Side effects of chemotherapy

Chemotherapy damages cells as they divide. This makes the drugs effective against cancer cells, which divide rapidly. However, some normal cells – such as hair follicles, blood cells, and cells inside the mouth or bowel – also divide rapidly. Side effects happen when chemotherapy damages these normal cells. Unlike cancer cells, normal cells can recover, so most side effects are temporary.

Nausea – You may feel sick for a few hours or days after each treatment. Not everyone feels sick, and you'll be given medicine to help prevent it. Some medicines may cause constipation; talk to your doctor about this.

Diarrhoea – You may have loose, watery stools and feel like you urgently need to go to the toilet. You may be given medicine to manage diarrhoea.

Hair loss – You may lose the hair from your head and other areas of the body (e.g. eyebrows, underarms and pubic area). Cold caps may prevent hair loss on your head in some cases (see page 54).

Swelling (oedema) – Some medicines used with chemotherapy drugs can cause excess fluid (fluid retention) to build up in the body. This can affect the arms and the trunk, but it usually gets better when treatment ends.

Changes to fertility – Chemotherapy can cause infertility in females and males. If you may want to have children in the future, it's essential that you talk to your cancer specialists about your options and ask for a referral to a fertility specialist before treatment starts (see page 57).

Heart problems – The risk is very small but chemotherapy can sometimes damage the heart muscle (cardiomyopathy). If necessary, your heart health may be checked before, during and after treatment. If you are at risk of heart damage, you may be offered other types of drugs.

Peripheral neuropathy – You might develop tingling or numbness in your hands or feet. This is called peripheral neuropathy (see page 53).

Other side effects – These include an increased infection risk, fatigue, mouth ulcers, constipation and memory changes (see page 54).

► See our *Understanding Chemotherapy* booklet.

Hormone therapy

Hormone therapy, also called endocrine therapy or hormone-blocking therapy, slows or stops the effect of oestrogen. It's used to treat breast cancer that is hormone receptor positive (see page 22). It's often used to lower the risk of the cancer coming back. It may also be used to reduce the risk of certain conditions, including some non-invasive breast conditions (see page 12), developing into invasive breast cancer.

There are different types of hormone therapy. The type used will depend on your age, type of breast cancer and if you have reached menopause.

Tamoxifen

Tamoxifen can be used at any age, whether you have been through menopause or not. You need to take a daily tablet for 5–10 years.

Side effects – In females, tamoxifen can cause menopausal symptoms (see page 57) like hot flushes and mood changes, although it doesn't bring on menopause. It may also cause changes in thinking and memory, and vaginal discharge. There is a very small risk of developing cancer of the uterus (endometrial cancer) if you have gone through menopause. Let your doctors know if you have any unusual vaginal bleeding. In males, side effects can include low sex drive (libido) and erection problems.

Tamoxifen increases the risk of blood clots. See a doctor immediately if you have swelling, soreness or warmth in an arm or leg, or a sudden shortness of breath or chest pain.

You are unlikely to have all of these side effects, and they usually improve with time. Your doctor and breast care nurse can help you to manage side effects. Tell your doctor if you take an antidepressant. Some types of antidepressant drugs may affect how well tamoxifen works.

Aromatase inhibitors

After menopause, the ovaries stop making oestrogen. However, both females and males make small amounts of oestrogen in body fat and the adrenal glands. Taking aromatase inhibitors will help reduce the amount of oestrogen in the body. This is important because oestrogen can cause some cancers to grow.

Aromatase inhibitors are mostly used if you have been through menopause, have had your ovaries removed or are male. They may also be used if you have not been through menopause but there is a high risk of the cancer coming back. If you have not gone through menopause, you may be advised to take an extra treatment to stop the ovaries from making oestrogen.

Side effects – Aromatase inhibitors can cause menopausal symptoms such as vaginal dryness, hot flushes and low mood (see page 57). These drugs may also cause itchiness, joint pain and stiffness, and weakening of the bones (osteoporosis). Your bone health will be monitored during treatment and you may be prescribed a drug to protect your bones. You should consider seeing a physiotherapist or exercise physiologist for an exercise plan. If you have arthritis, aromatase inhibitors may worsen joint stiffness and pain. Exercise or medicines from your doctor may help. Your doctor may also suggest changing to another type of aromatase inhibitor.

Ovarian suppression

If you have not been through menopause, drugs or surgery can stop the ovaries from producing oestrogen. This is called ovarian suppression. It may also be recommended as an additional treatment for people taking tamoxifen or for premenopausal women taking an aromatase inhibitor instead of tamoxifen.

Temporary ovarian suppression – A drug that stops the ovaries from producing oestrogen is given by injection. Side effects are similar to those of permanent menopause (see page 57). The drug may also help protect the ovaries during chemotherapy, so it's often given to women who want to preserve their fertility.

Permanent ovarian treatment – As an alternative to injections, some women choose to have their ovaries removed in a laparoscopic bilateral salpingo-oophorectomy, a small surgery where the ovaries and both tubes are removed.

Removing the ovaries will bring on permanent menopause. This means you will no longer be able to fall pregnant naturally.

Targeted therapy

Targeted therapy drugs attack specific features of cancer cells to stop the cancer growing and spreading. Different types of targeted therapy drugs are used for different types of breast cancer.

HER2-targeted agents

For early or locally advanced HER2+ breast cancer, the most common targeted therapy drug used is trastuzumab. It works by attaching itself to HER2+ breast cancer cells, destroying the cells or reducing their ability to divide and grow. It also encourages the body's own immune cells to help find and destroy cancer cells. It's usually used in combination with chemotherapy drugs for early breast cancer and it can increase the effect of the chemotherapy.

The drug can be given through a drip into a vein (infusion) or as an injection under the skin. The first infusion takes about 90 minutes (called

the loading dose). The following infusions each take 30–60 minutes. You will usually have a dose every 3 weeks, for up to 12 months. The first 4–6 doses are given while you are having chemotherapy treatment.

Side effects – Side effects are usually caused by the chemotherapy drugs, and often ease once chemotherapy finishes and you are having trastuzumab only. Side effects from trastuzumab are not common, but can include headache, fever and diarrhoea. In some cases, the drug can affect how the heart works, so you will have tests to check your heart function before and during treatment.

CDK inhibitors

These are a group of medicines that block cyclin-dependent kinases (CDKs), a group of proteins that help with regulating the cell cycle. They are used with hormone therapy to treat larger, high-risk ER+, HER2– breast cancers, for cancers involving several lymph nodes, or cancers at high risk of returning. They are used after surgery and chemotherapy.

Side effects – You may get nausea or diarrhoea, but this can be managed. Your blood count may be affected, so you will need regular blood tests.

PARP inhibitors

There are several new drugs for people who have inherited a BRCA mutation, or whose cancer has developed BRCA mutations. These are called poly (ADP-ribose) polymerase (PARP) inhibitors. Ask your doctor if they may be suitable for you.

► See our *Understanding Targeted Therapy* fact sheet.



Due to ongoing research, there are many new targeted therapy drugs for breast cancer. Your doctor will be able to tell you which ones may be suitable for you.

Immunotherapy

Immunotherapy is a treatment that uses the body's own immune system to fight cancer. A drug called pembrolizumab may be used for people with certain types of triple negative breast cancer. It is used together with a chemotherapy drug. It's given through a vein (intravenously), and treatments usually take about 30 minutes.

Side effects – These may be caused by the immunotherapy, the chemotherapy or both. Common side effects include a rash, fatigue, diarrhoea (which can be severe), breathlessness, joint pains, nerve problems, muscle weakness and dry eyes. It can cause inflammation in other organs, including the thyroid, pituitary gland, liver, kidneys and pancreas. It can also affect the adrenal gland, which can lead to low levels of certain hormones (e.g. cortisol).

Let your treatment team know if you notice these or any other side effects – some side effects can be life-threatening if they are left untreated. Most side effects can be managed if they are reported early. Sometimes, immunotherapy may need to be stopped or interrupted.

Side effects from immunotherapy can last for up to 12 months after the last dose was given. You should avoid falling pregnant during this time.

Occasionally, immunotherapy can cause adrenal failure and diabetes. It can also lead to thyroid conditions and low steroid levels. These complications need lifelong management.

► See our *Understanding Immunotherapy* fact sheet.

“It’s important to play an active role in your treatment because it’s your life and your body.” VIVIEN

Treatment for advanced breast cancer

Advanced breast cancer is different from locally advanced breast cancer (see page 21). Locally advanced breast cancer is cancer (larger than 5 cm) that has spread to tissue around the breast or to a large number of lymph nodes. Advanced breast cancer is cancer that has spread to more distant body parts. It is also called metastatic or secondary breast cancer.

Breast cancer can spread to many different parts of the body, but it is most likely to spread to the bones, liver, lungs or brain.

The treatment for advanced breast cancer varies from person to person. It will depend on the type of breast cancer and where in the body the cancer has spread.

The treatment for advanced breast cancer aims to control the spread of the cancer and relieve any symptoms you may develop. You may have one or more of the following treatments:

- hormone therapy (see pages 45–47)
- chemotherapy (see pages 43–44)
- targeted therapy (see pages 47–48)
- immunotherapy (see page 49).

In some cases, radiation therapy (see pages 40–42) may be used to reduce the size of the cancer and to relieve pain.

Surgery is not often used for advanced breast cancer, but it is occasionally used to remove some of the cancer that has spread to the bones, lungs, brain or liver.

While it's not possible to cure advanced breast cancer at this time, these treatments may improve quality of life for many months and, often, years.

For more information, see our other resources:

- *Living with Advanced Cancer*
- *Understanding Cancer Pain*
- *Understanding Secondary Bone Cancer*
- *Understanding Secondary Liver Cancer*
- *The Thing About Advanced Cancer* podcast series.

For more information about advanced breast cancer, visit Breast Cancer Network Australia at bcna.org.au and the MBC Hub from Metastatic Breast Cancer Action Australia at mbchub.au.

Key points about treating breast cancer

Choice of treatment

The treatments you are offered will depend on many factors, such as test results on the cancer, the stage and grade, whether it is hormone receptor positive, HER2+ or triple negative, and your age and general health.

Treatments before surgery

Before surgery, chemotherapy is usually offered to people with locally advanced, HER2+ or triple negative, or inflammatory breast cancer. This is called neoadjuvant chemotherapy. Some people also have hormone therapy, targeted therapy or immunotherapy before surgery.

Surgery

Early and locally advanced breast cancer is usually treated with surgery:

- breast-conserving surgery – removes only the tumour and a tissue margin
- mastectomy – removes the whole breast
- axillary surgery – removes some or all of the lymph nodes from the armpit to check for cancer; can be done through a sentinel lymph node biopsy (SLNB) or axillary lymph node dissection (ALND).

You may also have a breast reconstruction, which re-creates the shape of your breast during or after a mastectomy. You may choose to delay this surgery or not have a breast reconstruction.

Treatments after surgery

After surgery you may have chemotherapy, radiation therapy, hormone therapy for hormone receptor positive breast cancer, or targeted therapy for HER2+ breast cancer. You may also have immunotherapy.

Managing side effects

It will take time to recover from the physical and emotional changes caused by your treatment. Side effects can vary. Some people will experience just a few side effects, while others will have more.

Lymphoedema

Lymphoedema is the swelling (oedema) that develops when lymph fluid builds up in the tissues of part of the body, such as an arm, breast or the chest wall. When lymph nodes have been damaged or removed, lymph fluid may not be able to drain properly and builds up in the tissues.

Some breast cancer treatments may cause lymphoedema (e.g. surgery to remove lymph nodes and radiation therapy to the armpit). Lymphoedema can affect people at any time – during treatment or months or even years afterwards. Regular screening check-ups may be recommended for some people, so ask your treatment team if this is an option for you. Signs of lymphoedema in your arm to look for include the swelling of part of your arm or your whole arm; a feeling of tightness, heaviness or fullness in the fingers, wrist or arm; and aching in the affected area. These signs may begin gradually or come and go.

Some people experience pain, redness or fever, which can be caused by an infection called cellulitis in the area with lymphoedema. If you have any of these symptoms, see your doctor as soon as possible. Lymphoedema is easier to manage when diagnosed and treated early.

If you have lymphoedema, you can ask your doctor to refer you to a physiotherapist or lymphoedema practitioner. If you live in a rural

or regional area, you may need to travel to visit a lymphoedema practitioner. Some clinics provide telehealth consultations.

► See our *Understanding Lymphoedema* fact sheet.

Cording

Cording (axillary web syndrome) can develop weeks or months after breast surgery. It's caused by hardened lymph vessels and feels like a tight cord running from your armpit down the inside of the arm, sometimes to the palm of your hand. You may see and feel raised cord-like structures across your arm, chest or breast, which may limit how you move. Gentle exercises after surgery can help, as well as massage, physiotherapy or low-level laser treatment by a lymphoedema practitioner. Cording usually improves over a few months.

Nerve pain

Mastectomy and lymph node removal (see pages 30 and 32) can cause nerve pain in the arm or armpit, and mastectomy can cause nerve pain in the chest. This may be pins and needles, tingling or stabbing pain. It usually settles in a few weeks. Talk to your doctor if it's ongoing.

Some chemotherapy drugs can damage nerves in the hands and feet. This is called peripheral neuropathy. It can cause weakness, numbness, pins and needles, and burning or shooting pain. Symptoms usually improve over a few months, but they can be permanent.

Tell your healthcare team if you have any symptoms. Your doctor will help you manage pain and a physiotherapist and occupational therapist can give you exercises.

► See our *Understanding Peripheral Neuropathy and Cancer* fact sheet.

Hair loss

You may lose your hair during chemotherapy. Hair loss usually starts about 2 weeks after the first treatment. You can wear a wig, scarf, turban or hat or you might be comfortable leaving your head bare. You can try a few options and see what feels right. Generally, hair starts to grow back after treatment ends.

Some treatment centres offer cold caps (also called scalp cooling), which may prevent total head hair loss, but this depends on the chemotherapy drugs you have. These machines are not available at all treatment centres and are not suitable for everyone. Some people find them too uncomfortable and they are not usually covered by Medicare. Ask your treatment team if cold caps might be an option for you.

► See our *Understanding Hair Loss* fact sheet.

Thinking and memory changes

Some people notice changes in the way they think and remember information. This is known as cancer-related cognitive impairment, “chemo brain”, “cancer fog” or “brain fog”. The exact cause is unknown, but studies suggest it may be caused by the cancer or the cancer treatment, side effects such as fatigue, insomnia, pain and hormone changes, the surgical anaesthetic, and anxiety or depression.

For most people, these problems get better within the first year of finishing treatment. Others may have long-term effects. If you have severe or lasting changes to your thinking and memory, you can see a clinical psychologist or neuropsychologist for cognitive rehabilitation. Speak to your healthcare team about the services at your hospital.

► See our *Understanding Changes in Thinking and Memory* fact sheet and listen to our “Brain Fog and Cancer” podcast episode.

Breast prostheses

A breast prosthesis is a synthetic breast or part of a breast that's worn in a bra or attached to the body with adhesive. It helps give the appearance of a breast shape and can be used after breast surgery. You can use it some or all of the time or not at all.

Temporary prosthesis – In the first month or two after surgery, you may choose to wear a temporary light breast prosthesis called a soft form. This will be more comfortable next to the scar. A free bra and soft form are available through Breast Cancer Network Australia as part of the My Care Kit. To order a kit, speak to your breast care nurse or visit bcna.org.au/my-care-kit.

Permanent prosthesis – When you have recovered from treatment, you can be fitted for a permanent breast prosthesis. These are usually made from silicone gel and have the shape, feel and weight of natural breasts. It's best to see a trained fitter who can help you choose the right prosthesis. To find a fitter near you, call Cancer Connect on 13 11 20 or ask your breast care nurse for recommendations.

► See our resources on breast prostheses and reconstruction.

Feelings of loss and change

After a cancer diagnosis, it's common to feel a sense of grief or loss – for your health and wellbeing, your dreams or freedoms, your body, even what you can wear. Grief can feel like waves of sadness or being teary and usually settles over time. Sometimes you don't feel

grief until treatment is over. If you're concerned, call Cancer Connect on 13 11 20 to connect with a qualified professional and talk to someone about how you are feeling.

► See our *Emotions and Cancer* booklet.



The Look Good Feel Better program can help you to manage the appearance-related effects of cancer treatment and boost self-esteem. Call 1800 650 960 or visit lgfb.org.au.

Changes to body image and sex life

Breast cancer can affect how you feel about yourself (self-esteem) and make you feel self-conscious and less confident. These feelings are common; give yourself time to adapt. If you are finding it hard to adjust to changes, ask for support. Most cancer centres have counsellors who may be able to help.

Breast and chest appearance – It's normal to experience a range of emotions about the changes to your body. There's no rush to make a decision but there are several options that can help you feel you are taking back control. You may find having a breast reconstruction (see page 31) or wearing a breast prosthesis (see page 55) improves your confidence. Or you may prefer to “go flat” (see page 30). You may be able to keep your nipple and areola or have your nipple surgically reconstructed. You can also have an areola and nipple tattooed onto the breast or you may choose a decorative tattoo to cover scars.

Low libido – Breast cancer and its treatment (particularly hormone treatment) can also reduce your desire for sex (libido). You may miss the pleasure you felt from the breast or nipple being stroked or kissed during sex. This may be the case even if you have a reconstruction. If breast stimulation was important for arousal before surgery, you may need to explore other ways of becoming aroused. Some cancer treatment centres have sexual health clinics and other resources that may be able to help.

Vaginal dryness – Some treatments for breast cancer (particularly hormone therapy) can cause vaginal dryness, which can make penetrative

sex painful. You can try treatments such as lubricants and non-hormonal vaginal moisturisers. If this doesn't work, and sex is still painful, talk to your doctor about oestrogen cream.

- ▶ See our *Sex, Intimacy and Cancer* and *LGBTQI+ People and Cancer* booklets and listen to our “Sex and Cancer” podcast episode.

Menopause and infertility

Chemotherapy can cause your periods to stop for a short time or it may cause them to stop permanently (early menopause). Symptoms include hot flushes, trouble sleeping, vaginal dryness, reduced sex drive (libido), tiredness, dry skin, mood swings, weight gain and osteoporosis. Talk to your healthcare team about your symptoms. Several non-hormonal medicines work well for hot flushes. If you have vaginal dryness, talk to your doctor about trying the treatments mentioned above. In some states you can also ask for a referral to a menopause clinic.

If chemotherapy causes menopause, you will not be able to have children naturally. Talk to your doctor before treatment starts, as there may be ways to reduce the risk of early menopause or preserve your fertility.

If you find out you might not be able to fall pregnant, you may feel a great sense of loss. Talking to a counsellor or someone in a similar situation may help. For information about counselling services and support groups in your area, call Cancer Connect on 13 11 20.

- ▶ See our *Fertility and Cancer* booklet.



If you had treatment for hormone-sensitive breast cancer, avoid hormone-based contraception (“the pill”, implants or injections). Use condoms, diaphragms or non-hormone intrauterine devices (IUDs). If you had triple negative breast cancer, talk to your doctor.

Looking after yourself

Cancer can cause physical and emotional strain, so it's important to look after your wellbeing. Cancer Council has free booklets and programs to help you during and after treatment. Call Cancer Connect on 13 11 20 to find out more or visit your local Cancer Council website (see back cover).

Eating well – Cancer and treatment place extra demands on the body. Eating well before, during and after treatment can help you cope with side effects and reduce the risk of the cancer coming back. Side effects from treatment may make it harder to tolerate certain foods. If you are losing weight or finding it hard to eat, a dietitian can help you find what works for you. Alcohol can interact with some medicines, so check with your doctor if you want to drink during treatment. If you choose to drink alcohol, the guidelines recommend no more than 10 standard drinks a week and no more than 4 standard drinks on any one day.

► See our *Nutrition for People Living with Cancer* booklet.

How staying active can help

You may think it's important to rest during treatment and recovery, but research shows that exercise benefits most people at this time. There is also strong evidence that exercise can reduce the risk of breast cancer returning.

Being active can also help manage side effects such as tiredness,

improve circulation, boost mood, and speed up recovery.

Exercise reduces the risk of weight gain, which is associated with some treatments. An accredited exercise physiologist or physiotherapist can develop a program to suit you.

► See our *Exercise for People Living with Cancer* booklet.

Work and money – Cancer can change your financial situation, especially if you have extra medical expenses or need to stop working. Getting professional financial advice and talking to your employer can give you peace of mind. You can also check whether any financial assistance is available to you by asking a social worker at your hospital or treatment centre or by calling Cancer Connect on 13 11 20.

► See our *Cancer and Your Finances* and *Cancer, Work and You* booklets.

Relationships – Having cancer can affect your relationships with family, friends and colleagues in many different ways. Cancer is stressful, tiring and upsetting, and this may strain your relationships. The experience of cancer may also result in positive changes to your values, priorities or outlook on life. Give yourself time to adjust to what's happening, and do the same for those around you. It may help to discuss your feelings with other people.

► See our *Emotions and Cancer* booklet.

Complementary therapies – Complementary therapies are designed to be used alongside conventional medical treatments. Therapies such as massage, relaxation and acupuncture can increase your sense of control, decrease stress and anxiety, and improve your mood. Let your doctor know about any therapies you are using or thinking about trying, as some may not be safe or evidence-based.

► See our *Understanding Complementary Therapies* booklet.



Alternative therapies are therapies used instead of conventional medical treatments. They are unlikely to be scientifically tested, may prevent successful treatment of the cancer and can be harmful. Cancer Council does not recommend the use of alternative therapies as a cancer treatment.

Life after treatment

For most people, the cancer experience does not end on the last day of treatment. Life after cancer treatment can present its own challenges. You may have mixed feelings when treatment ends, and worry that every ache and pain means the cancer is coming back.

Some people say that they feel pressure to return to “normal life”. It is important to allow yourself time to adjust to the physical and emotional changes, and establish a new daily routine at your own pace. Your family and friends may also need time to adjust.

Cancer Connect can help you connect with other people who have had breast cancer and give you information about living well after cancer. Call 13 11 20 or visit cancerconnect.org.au.

► See our *Living Well After Cancer* booklet.

Dealing with feelings of sadness

If you have continued feelings of sadness, have trouble getting up in the morning or have lost motivation to do things that previously gave you pleasure, you may be experiencing depression. This is quite common among people who have had cancer.

Talk to your GP, because counselling or medication – even for a short time – may help. Some people can

get a Medicare rebate for sessions with a psychologist. Cancer Council may also run a counselling program in your area.

For information about coping with depression and anxiety, call Beyond Blue on 1300 22 4636 or visit beyondblue.org.au. For 24-hour crisis support, call Lifeline 13 11 14 or visit lifeline.org.au.

Follow-up appointments

After treatment ends, you will have regular appointments with your cancer specialist and GP to monitor your ongoing health. This is known as shared care. Your doctors will see how you are going on hormone therapy (if this is part of your ongoing treatment), help you to manage any long-term side effects such as lymphoedema, peripheral neuropathy or heart issues, and check that the cancer has not come back or spread.

During these check-ups, you will usually have a physical examination. You will also be able to discuss how you're feeling and mention any concerns you may have.

Check-ups after breast cancer treatment are likely to happen every 6–12 months for 2 years. They will become less frequent after that if you have no further problems.

You are likely to have a mammogram and, if needed, an ultrasound every year. You won't need a mammogram if you've had a double mastectomy. If there is a concern the cancer may have come back, you may have a bone scan and a CT, PET or MRI scan. After 5 years with no sign of cancer, women aged 40 and over can continue to have a free mammogram through the national breast cancer screening program.

When a follow-up appointment or test is approaching, many people find that they think more about the cancer and may feel anxious ("scanxiety"). Talk to your treatment team or call Cancer Connect on 13 11 20 if you are finding it hard to manage this anxiety.

Between follow-up appointments, let your doctor know immediately if you have any symptoms or health problems.

What if the cancer returns?

Sometimes, breast cancer does come back after treatment, which is called a recurrence. This is why regular check-ups are important. In most cases, early breast cancer will not come back (recur) after treatment. Although the risk is higher with locally advanced breast cancer, many people will not experience a recurrence.

There are some things that increase the risk that cancer may come back. These include if the cancer was large or the grade was high when first diagnosed, if it was found in the lymph nodes, or if the surgical margin was not clear. Your risk may also be increased if the cancer was hormone receptor negative or if adjuvant treatment (e.g. radiation therapy, chemotherapy, hormone therapy) was recommended after surgery but was not started or completed. This does not mean the cancer will definitely come back or spread.

Regularly looking at and feeling your breasts to know what is normal (being “breast aware”) can help find cancer in the treated or other breast.

If you have had a double mastectomy with or without a reconstruction, you should also regularly look at and feel your new shape and get to know your “new normal”. Tell your specialist, breast care nurse or GP if you notice any changes. Breast cancer can also return in other parts of the body, such as the bones, liver or lungs. Most symptoms will not be a recurrence, but if you notice any changes to your health, even years later, see your doctor and let them know that you have had breast cancer.

It is important to continue taking the drugs your doctor prescribes, even months or years after your treatment. Talk to your doctor before you stop taking any drugs, as they may be helping to stop the cancer from coming back.

Caring for someone with cancer

You may be reading this booklet because you are caring for someone with cancer. What this means for you will vary depending on the situation. Being a carer can bring a sense of satisfaction, but it can also be challenging and stressful.

It is important to look after your own physical and emotional wellbeing. Give yourself some time out and share your concerns with somebody neutral such as a counsellor or your doctor, or try calling Cancer Connect on 13 11 20. There is a wide range of support available to help you with the practical and emotional aspects of your caring role.

Support services – Support services such as Meals on Wheels, home help or visiting nurses can help you in your caring role. You can find local services, as well as information and resources, through the Carer Gateway. Call 1800 422 737 or visit carergateway.gov.au.

Support groups and programs – Many cancer support groups and cancer education programs are open to carers as well as to people with cancer. Support groups and programs offer the chance to share experiences and ways of coping.

Carers Australia – Carers Australia provides information and advocacy for carers. Visit carersaustralia.com.au.

Cancer Connect – Call 13 11 20 or visit cancerconnect.org.au to find out more about carers' services.

► See our *Caring for Someone with Cancer* booklet.

Seeking support

A cancer diagnosis can affect every aspect of your life. You will probably experience a range of emotions – fear, sadness, anxiety, anger and frustration are all common reactions. Cancer also often creates practical and financial issues.

There are many sources of support and information to help you, your family and carers navigate all stages of the cancer experience, including:

- information about cancer and its treatment
- access to benefits and programs to ease the financial impact of cancer treatment
- home care services, such as Meals on Wheels, visiting nurses and home help
- aids and appliances
- support groups and programs
- counselling services
- exercise programs.

The availability of services may vary depending on where you live, and some services will be free but others might have a cost.

To find good sources of support and information, you can talk to the social worker or breast care nurse at your hospital or treatment centre, or get in touch with Cancer Connect on 13 11 20.

“My family members don’t really understand what it’s like to have cancer thrown at you, but in my support group, I don’t have to explain.” SAM

Support from Cancer Council

Cancer Council offers a range of services to support people affected by cancer, their families and friends. Services may vary by location.

Cancer Connect by Cancer Council



Call 13 11 20 to talk to a qualified professional. We can answer questions you have about your situation and link you to local services (see inside back cover).

Information resources



Cancer Council produces booklets and fact sheets on more than 40 types of cancer, as well as treatments, emotional and practical issues, and recovery. Call 13 11 20 or visit cancerconnect.org.au.

Legal and financial support



If you need advice on legal or financial issues, we may be able to refer you to appropriate professionals. These services may be free for people who can't afford to pay. Financial assistance may also be available. To find out more, call Cancer Connect on 13 11 20.

Practical help



Cancer Connect can help you find ways to manage the practical impacts of cancer. This can include connecting you with services, such as accommodation and transport services.

Peer support services



You might find it helpful to share your thoughts and experiences with other people affected by cancer. We can link you with individuals or support groups by phone, in person, or online. Call 13 11 20 or visit cancercouncil.com.au/OC.

Useful websites

You can find many useful resources online, but not all websites are reliable. These websites are good sources of support and information.

Australian

Cancer Connect	cancerconnect.org.au
Cancer Council Online Community	cancercouncil.com.au/OC
Cancer Council podcasts	cancercouncil.com.au/podcasts
Guides to Best Cancer Care	cancer.org.au/cancercareguides
Australasian Lymphology Association	lymphoedema.org.au
Australasian Menopause Society	menopause.org.au
Australian Breast Device Registry	abdr.org.au
Breast Cancer Network Australia	bcna.org.au
Breast Cancer Trials	breastcancertrials.org.au
Breast Surgeons of Australia and NZ	breastsurganz.org
Cancer Australia	canceraustralia.gov.au
McGrath Foundation	mcgrathfoundation.com.au
National Breast Cancer Foundation	nbcf.org.au

International

American Cancer Society	cancer.org
Breast Cancer Now	breastcancer.org
Cancer Research UK	cancerresearchuk.org
Macmillan Cancer Support (UK)	macmillan.org.uk

Question checklist

Asking your doctor questions will help you make an informed choice. You may want to include some of the questions below in your own list.

Diagnosis

- What type of breast cancer do I have?
 - Has the cancer spread? If so, where has it spread? How fast is it growing?
 - Are the latest tests/treatments for this cancer available in this hospital?
 - Will a multidisciplinary team be involved in my care?
 - Are there clinical guidelines for this type of cancer?
-

Treatment

- What treatment do you recommend? What is the aim of the treatment?
 - Are there other treatment choices for me? If not, why not?
 - If I don't have the treatment, what should I expect?
 - How long do I have to make a decision?
 - I'm thinking of getting a second opinion. Can you recommend anyone?
 - How long will treatment take? Will I have to stay in hospital?
 - Are there any out-of-pocket expenses not covered by Medicare or my private health cover? Can the cost be reduced if I can't afford it?
 - How will we know if the treatment is working?
 - Are there any clinical trials or research studies I could join?
-

Side effects

- What are the risks and possible side effects of each treatment?
 - Will I have a lot of pain? What will be done about this?
 - Can I work, drive and do my normal activities while having treatment?
 - Will the treatment affect my sex life and fertility?
 - Should I change my diet or physical activity during or after treatment?
 - Are there any complementary therapies that might help me?
-

After treatment

- How often will I need check-ups after treatment?
 - If the cancer returns, how will I know? What treatments could I have?
-

Glossary

adjuvant therapy

A treatment given after the main treatment to lower the risk that the cancer will come back.

advanced cancer

Cancer that is unlikely to be cured. In most cases, the cancer has spread to other parts of the body (secondary or metastatic cancer). Treatment can often still control the cancer and manage symptoms.

areola

Coloured rim of tissue around the nipple.

aromatase inhibitors

Drugs that help prevent the growth of oestrogen-dependent cancer cells by reducing the amount of oestrogen in a postmenopausal woman's body.

atypical ductal hyperplasia (ADH)

An abnormal but non-cancerous condition of the cells in the lining of the milk ducts.

axilla

The armpit.

axillary dissection or surgery

The removal of some lymph nodes in the armpit to check whether cancer has spread.

axillary lymph nodes

Lymph nodes in and around the armpit.

axillary web syndrome

See cording.

biopsy

Removal of a sample of cells or tissue for examination under a microscope.

bone scan

A small amount of radioactive dye is injected into a vein. It collects in the bones and is detected by a scanning machine.

BRCA1 or BRCA2 mutation

A gene change that increases the risk of breast, ovarian or prostate cancer.

breast-conserving surgery

Surgery that removes a breast lump without removing the entire breast. Also called a lumpectomy or wide local excision.

breast form

See breast prosthesis.

breast oedema

Swelling caused by too much fluid in the breast tissue.

breast prosthesis (plural: prostheses)

An artificial breast worn inside a bra or attached to the body to re-create the shape of a natural breast. Also called a breast form.

breast reconstruction

Surgery to rebuild the breast shape after the breast, or part of it, is removed.

cancer-related cognitive impairment

Thinking and memory problems after treatment. Also called "chemo brain", cancer fog or brain fog.

cellulitis

An infection of the skin. It can happen after lymph glands have been removed.

chemotherapy

A cancer treatment that uses drugs to kill cancer cells or slow their growth.

clear margin

When the surrounding tissue removed during surgery does not contain cancer cells.

cold cap

A cap connected to a cooling system worn during chemotherapy to help prevent hair loss.

contrast enhanced mammogram (CEM)

Imaging that combines a 3D mammogram (tomosynthesis) with an injection of dye.

cording

Tight cords of tissue running down the inside of the arm. Also called axillary web syndrome.

CT scan

Computerised tomography scan. Uses x-rays to create a cross-sectional picture of the body.

diethylstilboestrol (DES)

A synthetic hormone drug identified as a cause of breast cancer.

DIEP flap reconstruction

A deep inferior epigastric perforator (DIEP) flap breast reconstruction. It uses skin and tissue from the stomach.

ductal carcinoma in situ (DCIS)

Abnormal cells that resemble cells in the breast ducts and may increase the risk of developing breast cancer.

ducts

The canals within the breast that pass milk from the lobules to the nipple.

early breast cancer

Cancer that has not spread beyond the breast or the axillary lymph nodes.

gene expression profile test

A test that gives information about the risk of the cancer coming back and whether chemotherapy will help. May also be called a molecular assay or genomic test.

genes

The units that determine how the body's cells behave, inherited from both parents.

genetic test

Aims to detect gene changes that may increase the risk of developing some cancers.

grade

A number that describes how similar cancer cells look to normal cells and how quickly they are likely to grow.

HER2

Human epidermal growth factor receptor 2. A type of protein involved in cell growth.

hormone receptors

Proteins in a cell that bind to specific hormones.

hormones

Chemicals in the body that send information between cells to control growth and reproduction.

hormone therapy

A treatment that blocks the body's natural hormones, which sometimes help cancer cells grow. Also called endocrine therapy or hormone-blocking therapy.

inflammatory breast cancer

Cancer that affects the lymphatic vessels in the skin of the breast.

invasive breast cancer

Cancer that has spread from the lining of the breast ducts or lobules into the breast tissue.

invasive ductal carcinoma (IDC)

Cancer that looks like cells in the milk ducts and has spread into the breast tissue.

invasive lobular carcinoma (ILC)

Cancer that looks like cells in the milk lobules and has spread into the breast tissue.

Klinefelter syndrome

A genetic syndrome in males who have 2 X chromosomes (XXY) instead of one (XY).

lobe

Part of the female breast that contains lobules.

lobular carcinoma in situ (LCIS)

Abnormal cells that resemble cells in the breast lobules and increase the risk of developing breast cancer.

lobules

The milk-producing glands in the breast.

locally advanced breast cancer

Cancer that is larger than 5 cm, has spread to tissues around the breast or has spread to a large number of lymph nodes.

lumpectomy

See breast-conserving surgery.

lymphatic system

A network of vessels, nodes and organs that removes excess fluid from tissues, absorbs fatty acids, transports fat and produces immune cells. Includes the bone marrow, spleen, thymus and lymph nodes.

lymph nodes

Small glands found in groups throughout the body. They help protect the body against infection. Also called lymph glands.

lymphoedema

Swelling caused by a build-up of lymph fluid.

lymph vessels

Thin tubes that carry clear fluid known as lymph all over the body.

mammogram

A low-dose x-ray of the breast tissue.

margin/surgical margin

An edge of tissue removed during surgery. A clear or negative margin means no cancer cells were found on the edge of the tissue. A positive margin means cancer cells were found on the edge of the tissue and further surgery is usually required.

mastectomy

Surgery to remove the whole breast.

menopause

When periods (menstruation) end.

metastasis (plural: metastases)

Cancer that has spread from a primary cancer in another part of the body. Also known as secondary or advanced cancer.

MRI scan

Magnetic resonance imaging scan. A scan using magnetic fields and radio waves to take detailed cross-sectional pictures of the body.

neoadjuvant treatment

A treatment given before another treatment

to help make the second treatment more successful.

nipple-sparing mastectomy

A type of mastectomy where the breast skin, nipple and areola are not removed.

non-invasive breast condition

Precancerous breast condition that is only in the ducts or lobules of the breast. Also called ductal or lobular carcinoma in situ (DCIS or LCIS).

oestrogen

A sex hormone made mainly by the ovaries. After menopause, the hormone is produced in the fat cells.

oncoplastic breast-conserving surgery

An operation in which the cancer is removed, and plastic surgery techniques are used to preserve the appearance of the breast as much as possible.

osteoporosis

Thinning and weakening of the bones that can lead to bone pain and fractures.

ovarian suppression

Methods to stop the ovaries from functioning.

Paget disease of the nipple

Cancer that develops in the breast ducts and spreads to the skin of the nipple and areola.

peripheral neuropathy

Weakness, numbness, tingling or pain, usually in the hands and feet, caused by damage to the nerves that are located away from the brain and spinal cord (the peripheral nerves).

PET scan

Positron emission tomography scan. A scan in which a person is injected with a small amount of radioactive solution to find cancerous areas.

progesterone

A sex hormone made mostly by the ovaries. It prepares the lining of the uterus for pregnancy.

radiation therapy

The use of targeted radiation to kill or damage cancer cells so they cannot grow, multiply or spread. Also called radiotherapy.

reconstructive surgeon (plastic surgeon)

A doctor who has had advanced surgical training in the restoration of skin and tissue to near-normal appearance and function.

recurrence

The return of a disease after a period of improvement (remission).

screening

An organised program to identify disease in people before any symptoms appear.

sentinel lymph node biopsy (SLNB)

Removal of the sentinel node (see below).

sentinel node

The first lymph node that breast cancer cells may spread to outside the breast.

seroma

A collection of fluid under a wound that may develop after surgery. It is not harmful but may need draining.

skin-sparing mastectomy

A mastectomy in which the whole skin of the breast, except the nipple and areola, is kept.

staging

Performing tests to work out how far a cancer has spread.

targeted therapy

Drugs that target specific cancer cell features to stop the cancer growing and spreading.

tomosynthesis

A digital x-ray mammogram machine that creates a three-dimensional image.

triple negative breast cancer

A breast cancer that does not have receptors for the hormones oestrogen or progesterone or for the growth factor HER2.

triple test

Collection of 3 tests to diagnose breast cancer. Includes physical examination, breast imaging (such as mammogram and ultrasound) and biopsy results.

tumour

A new or abnormal growth of tissue on or in the body.

ultrasound

A scan that uses soundwaves to create a picture of part of the body.

wide local excision

A surgical procedure to remove a cancer or tissue and some healthy tissue around it.

Can't find a word here?

For more cancer-related words, visit connect.cancer.org.au/glossary.

References

1. Cancer Australia, *Guidance for the Management of Early Breast Cancer: Recommendations and Practice Points*, Cancer Australia, 2020, canceraustralia.gov.au/publications-and-resources.
2. A Gennari et al., "ESMO Clinical Practice Guideline for the diagnosis, staging and treatment of patients with metastatic breast cancer", *Annals of Oncology*, vol. 32, iss. 12, 2021, pp. 1475–1495.
3. J Benitez Fuentes et al., "Global stage distribution of breast cancer at diagnosis: a systematic review and meta-analysis", *JAMA Oncology*, vol. 10, iss. 1, 2023, pp. 71–78.
4. Australian Institute of Health and Welfare (AIHW), *Cancer Data in Australia*, AIHW, Canberra, 2025, viewed 14 August 2026, available from aihw.gov.au/reports/cancer/cancer-data-in-australia.



How you can help

At Cancer Council, we're dedicated to improving cancer control. As well as funding millions of dollars in cancer research every year, we advocate for the highest quality care for cancer patients and their families. We create cancer-smart communities by educating people about cancer, its prevention and early detection. We offer a range of practical and support services for people and families affected by cancer. All these programs would not be possible without community support, great and small.

Join a Cancer Council event: Join one of our community fundraising events such as Daffodil Day, Australia's Biggest Morning Tea, Relay For Life, Girls' Night In and other Pink events, or hold your own fundraiser or become a volunteer.

Make a donation: Any gift, large or small, makes a meaningful contribution to our work in supporting people with cancer and their families now and in the future.

Buy Cancer Council sun protection products: Every purchase helps you prevent cancer and contribute financially to our goals.

Help us speak out for a cancer-smart community: We are a leading advocate for cancer prevention and improved patient services. You can help us speak out on important cancer issues and help us improve cancer awareness by living and promoting a cancer-smart lifestyle.

Join a research study: Cancer Council funds and carries out research investigating the causes, management, outcomes and impacts of different cancers. You may be able to join a study.

To find out more about how you, your family and friends can help, please call your local Cancer Council.

The right support.



Cancer is never one story.
Every diagnosis, every person,
and every path is different. You
don't have to figure it out alone.

Cancer Connect can help you find
the information and support that
are right for you.

You can explore trusted information
on our Cancer Connect website at your
own pace, or call 13 11 20 to talk to our
qualified professionals. We will listen to
your concerns, answer your questions,
and connect you with support.

Wherever you are on your path,
Cancer Connect can help make the
next step feel clearer.

Right when you need it.



Call 13 11 20



Visit cancerconnect.org.au



If you need information in a language other than English,
an interpreting service is available. Call 131 450.



If you are deaf, or have a hearing or speech impairment,
you can contact us through the National Relay Service.
accesshub.gov.au

*Cancer Council services and programs vary in each area.
13 11 20 is charged at a local call rate throughout Australia (except from mobiles).*



For information & support,
call **Cancer Connect 13 11 20**
or visit **cancerconnect.org.au**

Cancer Council ACT
actcancer.org

Cancer Council Queensland
cancerqld.org.au

Cancer Council Victoria
cancervic.org.au

Cancer Council NSW
cancercouncil.com.au

Cancer Council SA
cancersa.org.au

Cancer Council WA
cancerwa.asn.au

Cancer Council NT
cancer.org.au/nt

Cancer Council Tasmania
cancer.org.au/tas

Cancer Council Australia
cancer.org.au

*This booklet is funded through the generosity of the people of Australia.
To support Cancer Council, call 13 11 20 or visit cancerconnect.org.au.*